INDEX TO SCIENTIFIC NAMES
HERPETOLOGICA
Volume 48 December 1992

Acanthophis antarcticus 146
Aextoxicum punctatum 116
Agama stellio 339
Agave 108
Ailuronyx 126, 129, 132
  seychellensis 125, 127, 131
Alligator mississippiensis 344
Alsodes 115
Amblyrhynchus cristatus 57
Ambystoma maculatum 284
  mexicanum 145
  tigrinum 284
Amolops larutensis 434–435, 437
Amphibolurus 145
Amphiuma 327
Andropogon spp. 307
Anguis 255, 258, 260–261
  fragilis 247, 343
Anniella 249–261
  pulchra 246–247
Anolis 366
  carolinensis 339
Anomalopus leuckarti 345
Antillophis andrea 173
Aristelliger hechti 131
  praesignis 131
Aristida spp. 307
Arrhyton ainictum 172, 174, 176–177
  bivittatum 171, 177
  callilaemum 176
  dolichura 168–172, 174, 176–177
  exiguum 175–176
  fulvum 177
  funereum 176
  landoi 168, 171–174, 176–177
  polylepis 176
  procerum 168, **174**–176
  redimitum 168–171, 177
  supernum 168, **171**–176
  taeniatum 168–172, 174, 176–177
  tanyplectum 170–172, 174, 176–177
  vittatum 168–172, 174, 176–177
Atractus 139
Aturia 85
Austrelaps superbus 343
Barygenys flavigularis 185, 190
Billbergia spp. 41
Boa constrictor 341
Bothrochilus perthensis 145–146
Bothrops 270
  asper 135, 139
  atrox 139
  colombiensis 139
  microphtalmus 135
Bufo americanus 215–216, 383–385, 387–388
  cognatus 41, 211
  hemiophrys 211, 383, 385, 387–389
  japonicus 284
  marinus 145
  microscaphus californicus 383–388
  microcaphus microscaphus 383–388
  punctatus 389
  woodhousii 383, 388
  woodhousii australis 388
  woodhousii woodhousii 388
Bungarus caeruleus 82
Buteo buteo 298
Caretta 6, 14
  caretta 1, 31, 38, 396
Carettochelys insculpta 145
Carinascincus greeni 345
  metallicum 343
  ocellatum 343
  pretiosum 343
Casarea dussumieri 146
Cassia spp. 41
Chalcides 340
  ocellatus tiligugu 339, 341
  tridactylus 343
Chamaeleo lateralis 343
Chelonia depressa 22
  mydas 6, 14, 21–22, 31, 38, 203, 269, 396
Chelydra 267
  serpentina 193, 200, 202, 205, 265–266, 268, 270, 344
Chiromantis 210
Chironius 137
  grandisquamis 138–139
  melas 138–139
  monticola 138
  multiventris 138
  multiventris cochranae 138
  schlueteri 138–139
  scurrulus 138
  spp. 140
Chitulia 85, 90–95
Chrysemys picta 193, 197, 200, 202, 344
Cistus ladanifer 294
Claireascincus entrecasteauxi 343
Claudius 267–268, 270
  angustatus 265–266, 269
Clelia 140
  scytalina 138
Cnemidophorus angusticeps 417–424
  cozumela 417–422
  cozumela cozumela 417–419, 421–424
  cozumela maslini 417–419, 421–424
  deppei 417–420, 422, 424
  guttatus 417–420, 422, 424
  lemniscatus 418–419
  motaguae 417–422, 424
  rodecki 417–422, 424
  sexlineatus 311, 330–333, 335, 339–341, 417
  sp. G 417–418, 420–421, 424

Colorhogia redimita 169
Coluber 448, 452
constrictor 220
Conchoderma virgata 89
Coniophanes fissidens 448–453
joanae 135
Conolophus pallidus 57
subcristatus 203
Conophis lineatus 136, 138
Cophixalus 184, 191
concinnus 186–187, 189–190
hosmeri 186–187, 189–190
ornatus 186–187, 189–190
pansus 185, 190
sp. B 186–187, 189–190
Copiula fistulans 184, 186–190
Cricosaura 107
typica 106
Crocodylus johnstoni 344
porosus 220, 344
siamensis 220
Crotalus 78
Ctenophorus reticulatus 345
Ctenotus robustus 345
Cupriguanus scapulatus 339–340
Cyclura nubila 203
Cynops 292
cyanurus 287–291
ensicauda 178, 181–182, 414
orientalis 287–291
pyrrhogaster 141
Darlingtonia haetiana 175
Delma fraseri 345
Delonix regia 41
Dendrophidion 140, 448
Dermatemys 267
mawii 265–266, 268–269
Dermochelys coriacea 22
Desmognathus 317
ochrophaeus 328
quadramaculatus 313–316
Diplodactylus strophurus 345
Dipsas pratti 135
Disteira 85
bituberculatus 86
Distira pachycercos 89
Drepanoides anomalus 140
Drosophila 126, 374
Drymarchon 139
Drymobius 448
Duellmanohyla chamulae 153–154, **163**–166
ignicolor 154, 164–166
lythrodes 154, 164–165
rufioculis 154, 164–165
salvavida 154, 164–165
schmidtorum 154, 164–166
soralia 154, 164–165
uranochroa 154, 164–165
Echinotriton 178
Egernia cunninghami 343
striolata 343
whitei 343
Eleutherodactylus 190–191, 366
amadeus 354, 356
antillensis 216
auriculatoides 355
auriculatus 352
bakeri 351, 354–357
buckleyi 347, 350
***cacao* 347**–350
coqui 47–49, 51–55, 216
***corona* 351**–357
cryophilius 350
curtipes 347, 350
diastema 47
dolomedes 351–354, **356**–357
eunaster 354, 356
fowleri 355
glanduliferoides 356
glaphycompus 356
gryllus 352, 355
haitianus 355
heminota 353–356
hypostenor 355
jamaicensis 355
jasperi 352, 355–356
johnstonei 40–41, 47–48
lamprotes 353, 355
latidiscus 347
limbatus 355
ruthae 355
semipalmatus 356
sisyphodemus 355
sulculus 347–348, 350
supernatis 347, 350
thorectes 355–356
thymalopsoides 347, 350
vertebralis 347, 350
Elgaria multicarinata 275
Emoia 149
Enulius 448
Eretmochelys imbricata 22, 396
Eucryphia cordifolia 116
Eulamprus "quoyii" 343
tympanum 390–394
Eulepis platynotum 345
Eumeces 305
fasciatus 229, 234, 307, 310–311
inexpectatus 307, 309–311
laticeps 230, 234, 307
Euphlyctis 437
Eupsophus 190
calcaratus 115, 117–119
contulmoensis 115
emiliopugini 115, 117–119
insularis 115
migueli 115
roseus 115, 117–118
vertebralis 115–119
vittatus 115
Eurycea 284
Falco tinnunculus 298
Gaigeia 97–98, 106
dontomasi 97, 105–107
gaigeae 108
radula 97, 105–106, 109
sylvaticum 97, 106
Geckolepis 129, 132
typica 124, 131
Gehyra mutilata 132

Gekko gecko 127, 129, 131
Gerrhosaurus nigrolineatus 229–234
Gevuina avellana 116
Gonocephalus grandis 120–123
miotympamum 120–123
spinipes 120–121
Gopherus polyphemus 310, 402
Graptemys 269
Halimium viscosum 294
Hannemania hylae 217
Harrisoniascincus coventryi 344–345
Heliconia 441, 444
Helicops leopardinus 137
Hemachatus 424, 433
Hemiergis decresiensis 344–345
Hinulia quadridigitatum 344
Hoplodactylus maculatus 343
Hydromantes 149
Hydrophis belcheri 89, 93, 95, 97
bituberculatus 85–86, 88–97
brooki 91
brookii 91–92, 97
caerulescens 92, 94, 96
cantoris 91
coggeri 86, 93, 97
cyanocinctus 86, 90, 94, 96–97, 343
czeblukovi 86, 93, 97
elegans 91
fasciatus 90–91, 97
geometricus 86, 93, 95
gracilis 91
inornatus 93, 95–97
klossi 91–92, 97
lamberti 86, 93, 95–97
lapemoides 92, 94, 96
macdowelli 86, 91, 93, 97
mamillaris 93, 95, 97
mcdowelli 91
melanocephalus 94, 97
melanosoma 91–92, 97
obscurus 91–92, 97
ornatus 90, 92, 94, 96–97
pacificus 86, 94, 97
parviceps 91–92, 96
spiralis 94, 97
stricticollis 93, 95–97
torquatus 93, 95–97
vorisi 86, 91, 93, 97
Hydrosaurus 57
Hyla 149
alleei 165
allenorum 439–444, 446–447
arenicolor 210–212, 214–217
avivoca 379
bokermanni 439, 445–447
brevifrons 439, 444–447
calcarata 439, 442–443, 447
chrysoscelis 244, 373–374, 376–381
cinerea 49, 243–244, 379
colymba 162
ebraccata 54, 374
erythromma 157, 165–166
euthysanota 161, 165
fasciata 439–443
geographica 439, 442–443
gratiosa 379
hutchinsi 439, 443
koechlini 439, 444–447
lanciformis 443, 447
legleri 156, 161, 165–166
leonhard-schultzei 165
leonhardschultzei 164
leucophyllata 439, 444
lythrodes 165
macrotympanum 161, 165
microcephala 54, 374, 439, 444
microps 439, 444–447
milleri 165
miotympanum 154, 162–163
mixomaculata 165
panchoi 163
parviceps 439, 444–447
phlebodes 374
rozellae 165
rufioculis 161, 165
salvadorensis 156, 161, 163, 165–166
salvavida 163, 165
schmidtorum 163
schubarti 439, 444
soralia 165
spinipollex 161, 163, 165
squirella 379
subocularis 445–447
uranochroa 161, 163, 165
versicolor 373–374, 376–381
Hylophorbus 185
rufescens 186–188, 190
Hymenocallis caribaea 41
Hynobius hidamontanus 286
kimurae 408–414
lichenatus 408–414
nigrescens 279–280, 283–286
retardatus 280, 285–286, 408–414
Hyperolius 210
marmoratus 48, 236–237, 243–245
Hypsilurus 122
godeffroyi 121
Imantodes gemnistratus 135
Japalura brevipes 123
Kinosternon leucostomum 265–269
Lacerta 330, 339
lepida 340
vivipara 340, 343, 393–394
Lachesis muta stenophrys 137
Lampropeltis 135
getula 341
getulus 220
Lanius excubitor 298
senator 298
Leioselasma 85, 90–91, 94
bituberculatus 86
Lepidochelys 14
kempi 22
olivacea 22
Lepidophyma 99–102, 276
dontomasi 97–98, 105–110
flavimaculatum 105, 107–108
gaigeae 97–98, 105–110
micropholis 105, 107
occulor 107–109

pajapanensis 105, 107
radula 97, 106–110
smithii 105, 107–109
sylvaticum 105–107, 109
tarascae 105, 108
tuxtlae 105, 107–108
Leptobrachium hasselti 437
hendricksoni 434–435, 437
Leptolalax pelodytoides 437
Liasis perthensis 145
Liolaemus austromendocinus 339–340
gravenhorti 343
Liophis 139
cobella 137
miliaris chrysostomus 137
poecilogyrus 138
ssp. 137
typhlus forsteri 138
undulatus 138
Lygodactylus picturatus 341
Lygosoma quadridigitatum 344
Malpolon monspessulanus 298
Mantidactylus 190
Marmorosphanx tricolor 345
Menetia greyii 345
Microcephalophis 85, 91
Microhyla butleri 434–438
heymonsi 435–437
Micrurus annelatus 138
dissoleucus dissoleucus 135
hemprichii 140
lemniscatus 137
nigrocinctus 449
nigrocinctus mosquitensis 135
spp. 453
spurelli 135
Morelia amethistina 145
Morethia obscura 345
Naja atra 425–427, 430–433
cf. *atra* 425–427, 431–432
kaouthia 69, 78–79, 84, 425–432
katiensis 432
mossambica 432
naja 69, 76–83, 424–432
naja bombaya 69, 81
naja gangetica 69, 81
naja indusi 69, 80
naja kaouthia 69–70, 77, 426, 432
naja karachiensis 69, 81
naja madrasiensis 69, 80
naja naja 69, 76, 80
naja oxiana 69, 76–77
nigricollis 432–433
oxiana 69, 76–79, 84, 424–432
pallida 432
philippinensis 69, 424–432
samarensis 69, 425–429, 431
sputatrix 69, 424–429, 431–433
sumatrana 69, 425–428, 431, 433
Natriciteres 448
Natrix natrix 80
Necturus alabamensis 318
beyeri 318–319, 321–322, 329
lewisi 318–320, 322–329
maculosus 59, 318, 321, 326–329
punctatus 318–319, 322, 328–329
Neobatrachus sudelli 146
Nerodia 448
erythrogaster 60, 62–63, 65–66
fasciata 65
fasciata confluens 60, 62–63, 65
harteri 60–62, 65–67
harteri harteri 60–63, 65–66
harteri paucimaculata 60–63, 65–67
paucimaculata 60, 67
rhombifera 60, 62–63, 65–66
taxispilota 60, 62–63, 65
Niceforonia 111
Ninia 140
Nobella 111
Nolina 108
Nothofagus 115
Notophthalmus 181
viridescens 216, 279, 283, 285
Ophisaurus 258
apodus 247, 255, 260–261
Oreophryne 185
biroi group sp. 1 186–187, 189–190
Oxyrhopus 140
melanogenys 137
Oxyuranus scutellatus 345
Pachydactylus scutatus 132
Perochirus spp. 132
Pheretima hilgendorfi 280
Philautus asper 434–437
leucorhinus 55
Philodryas olfersi 138
Phrynomantis robusta 186–188, 190
slateri 186–190
Phrynopus cophites 112
***kempffi* 111**–114
laplacai 111–112, 114
montium 112
peruanus 112, 114
wettsteini 112
Phyllomedusa 210
Phymaturus palluma 339–340
Physignathus lesueurii 345
Pinus palustris 307
Pituophis melanoleucus 220
Platylepas ophiophilus 89
Plectrohyla 163–164
Plethodon cinereus 288
jordani 216
Pleurodeles 327
Podarcis muralis 220–221, 223–226
Pogonia barbata 145, 345
vitticeps 345
Polypedates 434
leucomystax 435–437
leucomystax linki 437
Porcellio scaber 280
Pristidactylus scapulatus 339
Proteus anguinus 318–322, 327–329
Psammodromus algirus 293–294, 298–299
Psammophis 448
Pseudechis porphyriacus 345
Pseudemoia spenceri 344–345

Pseudemydura umbrina 146
Pseudonaja textilis 146
Pseudophryne 55
Pseudopus apodus 247
Pseudosalamandra 414
Ptenopus 55
Ptychohyla adipoventris 161, 164–165
bogerti 165
chamulae 155–162, 165
erythromma 154, 162, 164–165
euthysanota 154–155, 160–166
euthysanota euthysanota 161, 165
euthysanota macrotympanum 161, 165
ignicolor 155, 161–162, 165
legleri 154, 162, 164–165
leonhard-schultzei 165
leonhardschultzei 154–155, 161–164
lythrodes 155, 162
macrotympanum 153–155, 161–166
merazi 154–155, 160–166
panchoi 154–155, 160–166
rozellae 165
rufioculis 155, 162
salvadorensis 154, 162, 164–165
salvavida 155, 162
sanctaecrucis 153–154, **155**–160, 163–166
schmidtorum 155, 161–163, 165
schmidtorum chamulae 161, 165
schmidtorum schmidtorum 161, 165
soralia 155, 162
spinipollex 154–155, 159–162, 164–165
uranochroa 155, 162
Quercus geminata 307
laevis 307
rotundifolia 294
Rana 190
blythii 434–437
chalconota 435–436
clamitans 374, 379
glandulosa 435–436
grunniens 437
hosii 435–436
macrodon 434, 437
magna 434, 437
pipiens 57–58, 215
sphenocephala 288–289
virgatipes 49, 374, 379
Ranidella parinsignifera 374, 379
signifera 374, 379
Rhacophorus 434
promianus 435–437
Rhadinaea 135, 448
occipitalis 138
Rheobatrachus 145
Rheodytes leucops 145
Rhinobothryum lentiginosum 137
Salamandrella 414
keyserlingii 408–413
Sancevieria spp. 41
Saproscincus mustelina 345
Satobius 408–409, 414
Sauromalus obesus 57–58
Sceloporus 225, 301
graciosus 393
jarrovi 394
poinsetti 394
undulatus 275, 393
woodi 393
Schaphiodontophis 448
Serenoa repens 307
Sibynophis 448
Sphaerodactylus argus 359
armasi 359
armstrongi 366
armstrongi hypsinephes 365
bromeliarum 359
celicara 359
cricoderus 358, 360, **362**–366
cryphius 366
darlingtoni 366
docimus 359
elasmorhynchus 366
elegans 359
gaigeae 366
goniorhynchus 366
intermedius 359
klauberi 366
micropithecus 366
nicholsi 366
nigropunctatus 358–359
notatus 359, 362
nycteropus 366
oliveri 359
omoglaux 366
parthenopion 361, 366
perissodactylius 366
ramsdeni 358–359, 361–367
richardsoni 131
roosevelti 124–132
ruibali 359
scaber 359
schwartzi 358–**359**, 360–366
sp. nov. 2 366
spp. 125
streptophorus 366
torrei 358
townsendi 366
Sphenodon 107
punctatus 344
Sphenophryne adelphi 186–187, 189–191
fryi 186–187, 189, 191
Fryi 190
gracilipes 191
pluvialis 186–187, 189–191
robusta 186–187, 189–191
schlaginhaufeni 184, 186–188, 190
sp. B 184, 186–190
Staurotypus 268, 270
triporcatus 265–267, 269
Stenorrhina 140
Sternotherus odoratus 193, 197, 200
Storeria occipitomaculata 393
Tachyglossa 344
Taricha 181
torosa 285
Telmatobius 111
Tenebrio 272

Teratoscincus 132
scincus 131
Terrapene carolina 6
Thamnodynastes gambotensis 138–139
Thamnophis 61, 448, 452
cyrtopsis 217
sirtalis 343
Thecadactylus rapicauda 131
Tilandsia 362
Tiliqua rugosa 301–302
Tilqua rugosa 301
Trachemys 267, 269
scripta 265–266, 268, 270
Trachydosaurus rugosus 301–303, 305
Trionyx triunguis 203
Triturus 149, 181
Topidurus 340–341
torquatus 339
Tylototriton 327
andersoni 178–179, 181–182
Tympanocryptis lineata 345
Uta stansburiana 393
Varanus gilleni 278
varius 146
Vibrissaphora 437
Vipera aspis 344
berus 393
russelli 82
Xantusia 97, 106–107
henshawi 109
riversiana 271–278
vigilis 108, 276
Xenobatrachus 185
mehelyi 184, 186–188, 190
Xenopholis scalaris 138
Xenopus 149
laevis 215

TOPICAL INDEX
HERPETOLOGICA
Volume 48 December 1992

AGGRESSION, *Xantusia riversiana* (thermal dependence of locomotion and aggression) 271–279.

AGONISTIC BEHAVIOR, *Eleutherodactylus johnstonei* 40–49.

ALLOZYMES, *Salamandrella keyserlingii*, *Hynobius kimurae*, *H. lichenatus*, and *H. retardatus* (relationships and taxonomic problems) 408–416; *Tylototriton andersoni* (3 islands of the Ryuku Archipelago) 178–184.

ANNUAL ACTIVITY, *Eumeces fasciatus* and *E. inexpectatus* (in burned Florida scrub and sandhill habitat) 307–312.

BEHAVIOR, *Caretta caretta* (nesting) 1–30; *Caretta caretta* and *Chelonia mydas* (effects of artificial lighting on nesting) 31–39; *Cynops cyanurus* and *C. orientalis* (foraging behavior compared) 287–292; *Eleutherodactylus coqui* (aggressive calls) 49–56; *Eleutherodactylus johnstonei* (male mating behavior on Barbados) 40–49; *Gerrhosaurus nigrolineatus* (tongue-flicking response of males to cloacal stimuli) 229–236; *Hyla chrysoscelis* and *H. versicolor* (calling sites in sympatric choruses in Missouri) 373–382; *Hynobius nigrescens* (wandering) 279–287; *Hyperolius marmoratus* (male behavior and mating success) 236–246; *Psammodromus algirus* (basking around shrubs) 293–300; *Xantusia riversiana* (thermal dependence of locomotion and aggression) 271–279.

CLADOGRAM, *Nerodia* (6 spp.) 65; *Ptychohyla* and *Duellmanohyla* 164.

CLOACA, proteid salamanders (anatomy) 318–329.

CLOACAL GLANDS, *Gerrhosaurus nigrolineatus* (appear to produce sex pheromones) 229–236; proteid salamanders (anatomy) 318–329.

DISTANCE-WAGNER TREE, *Salamandrella keyserlingii*, *Hynobius kimurae*, *H. lichenatus*, and *H. retardatus* 412.

DISTRIBUTION MAP, *Arrhyton procernum* and *A. supernum* 175; *Naja kaouthia*, *N. naja*, and *N. oxiana* 79; *Naja* (9 Asiatic spp.) 425; *Sphaerodactylus ramsdeni* group (E Cuba) 362.

ECOLOGY, *Eumeces fasciatus* and *E. inexpectatus* (related to burning in Florida scrub and sandhill habitat) 307–312.

ECOMORPHS, *Sphaerodactylus* (mesic forest and xeric forest) 358–367.

EGG LAYING, *Caretta caretta* 1–30.

EGG SURVIVAL, *Podarcis muralis* (effects of incubation temperature on duration of incubation, embryo survival, and hatchling performance) 220–228.

EMBRYOLOGY, squamates (orientation of embryo on yolk mass) 343–346.

ETHOGRAM, *Caretta caretta* (nesting female) 1–30.

EVOLUTION, *Cnemidophorus cozumela* species group (matrilineal history and age) 417–424.

FANG STRUCTURE, *Naja* (related to spitting in 9 Asiatic cobras) 424–434.

FORAGING, *Cynops cyanurus* and *C. orientalis* (search behavior compared) 287–292.

GEOGRAPHIC VARIATION, *Naja* (fang structure of some Asiatic spp.) 424–434.

GROWTH RATE, *Podarcis muralis* (affected by incubation temperature) 220–228.

HABITAT, *Gopherus polyphemus* (estimating population size from number of burrows) 402–407.

HATCHLINGS, *Caretta caretta* (emergence pattern on Cephalonia) 396–401; *Podarcis muralis* (size affected by incubation temperature) 220–228.

HELIOTHERMY, *Psammodromus algirus* 293–300.

HEMIPENES, *Hydrophis bituberculatus* 89.

HISTOLOGY, *Chelydra serpentina* (testicular cycle in Wisconsin) 193–201; *Gerrhosaurus nigrolineatus* (cloacal anatomy) 229–236; *Sphaerodactylus roosevelti* skin 124–133.

HOME RANGE, *Tiliqua rugosa* (in S Australia) 301–306.

HYDRATION, *Hyla arenicolor* (effects of body temperature and hydration on metabolism) 210–219.

INCUBATION, *Podarcis muralis* (effects of incubation temperature on duration of incubation, embryo survival, and hatchling performance) 220–228.

INTERSPECIFIC INTERACTIONS, *Hyla chrysoscelis* and *H. versicolor* (calling sites in sympatric choruses in Missouri) 373–382.

KARYOTYPES, *Gonocephalus grandis* and *G. miotympanum* 122; *Lepidophyma gaigeae* and *L. dontomasi* 106; Malaysian frogs (11 spp.) 434–438; microhylid frogs (16 spp. in 7 genera from New Guinea and Australia) 184–192.

LARVAL DEVELOPMENT, *Desmognathus quadramaculatus* (in NE Georgia) 313–317.

LIMBLESSNESS, *Anniella pulchra* (lateral undulation, concertina, and slide-pushing locomotion) 246–262.

LITTER SIZE, *Eulamprus tympanum* (related to female size) 390–395.

LOCOMOTION, *Anniella pulchra* (traversing plane surfaces, fields of pins and nails, and channels) 246–262; *Caretta caretta* gait 1–30; *Xantusia riversiana* (effect of temperature on sprint speed) 271–279.

METABOLISM, *Hyla arenicolor* (effects of body temperature and hydration on metabolism) 210–219.

MITOCHONDRIAL DNA, *Cnemidophorus cozumela* species group (matrilineal history and age) 417–424; *Nerodia* (evolution and speciation in 6 spp.) 60–68.

MORPHOLOGY, *Naja* (fang structure related to spitting in 9 Asiatic cobras) 424–434.

NESTING, *Caretta caretta* (behavior) 1–30; *Caretta caretta* (emergence pattern of hatchlings on Ceph-

alonia) 396–401; *Caretta caretta* and *Chelonia mydas* (effects of artificial lighting on nesting behavior) 31–39.
OFFSPRING SIZE, *Eulamprus tympanum* (related to female size) 390–395.
OXYGEN CONSUMPTION, *Xantusia riversiana* (following burst sprinting) 271–279.
PARTHENOGENESIS, *Cnemidophorus cozumela* species group (matrilineal history and age) 417–424.
PHENOGRAM, 6 spp. of *Nerodia* (based on mtDNA) 63.
PHEROMONES, *Gerrhosaurus nigrolineatus* (experimental evidence for involvement in sex discrimination) 229–236.
PHOTOGRAPHS, *Arrhyton dolichura* and *A. procernum* 174; *Arrhyton landoi, A. supernum, A. taeniatum,* and *A. vittatum* 172; *Chelydra serpentina* (spermatogenesis) 196, 198; *Cnemidophorus sexlineatus* (sperm ultrastructure) 332, 334, 336–337; *Eleutherodactylus corona* and *E. dolomedes* (new spp. from Haiti) 352, type locality 354; *Gerrhosaurus nigrolineatus* (cloacal histology) 229–236; *Hydrophis bituberculatus* 88; *Hydrophis* palatine bones (subgenera *Hydrophis, Leioselasma, Chitulia*) 90; *Hynobius nigrescens* (skin histology) 285; *Lepidophyma gaigeae* and *L. dontomasi* 98, 108, *L. radula* 108; *Lepidophyma gaigeae* and *L. dontomasi* habitats 98; *Phrynops laplacai* and *P. kempffi* (new sp. from Bolivia) 112; proteid salamanders (cloacal anatomy) 320–321, 324–326; *Ptychohyla sanctaecrucis* (new sp. from Guatemala) 156; *Sphaerodactylus cricoderus, S. schwartzi,* and *S. ramsdeni*) 360; *Sphaerodactylus roosevelti* 125; *Sphaerodactylus roosevelti* (skin histology) 130.
PHOTOPOLLUTION, *Caretta caretta* and *Chelonia mydas* (nesting beaches) 31–39.
PHOTOTAXIS, *Caretta caretta* 1–30.
PHYLOGENY, *Salamandrella keyserlingii, Hynobius kimurae, H. lichenatus,* and *H. retardatus* (relationships and taxonomic problems) 408–416; squamates (orientation of embryo on yolk mass) 343–346.
PHYLOGRAM, *Cnemidophorus cozumela* species group 420; *Nerodia* (6 spp.) 65.
PHYSIOLOGICAL ECOLOGY, *Caretta caretta* (correlation between cooling sand and emergence of hatchlings) 396–401; *Hyla arenicolor* (effects of body temperature and hydration on metabolism) 210–219; *Podarcis muralis* (effects of incubation temperature on duration of incubation, embryo survival, and hatchling performance) 220–228; *Rana pipiens* and *Sauromalus obesus* (heating and cooling rates compared) 57–60; turtle nests (measuring water potential by thermocouple psychometry) 202–209.
POPULATION ESTIMATES, *Gopherus polyphemus* (method of counting burrows and applying a "correction factor" analyzed) 402–407.
PREY AVAILABILITY, *Cynops cyanurus* and *C. orientalis* (effects on search behavior) 287–292.
PROLACTIN, *Hynobius nigrescens* (effect on wandering behavior and skin) 279–287.
REPRODUCTION, *Bufo microscaphus* (calling behavior and advertisement and release calls) 383–389; *Caretta caretta* (nesting behavior) 1–30; *Caretta caretta* and *Chelonia mydas* (disruption of nesting by beach lighting) 31–39; *Chelydra serpentina* (testicular cycle in Wisconsin) 193–201; *Eleutherodactylus johnstonei* (alternative male reproductive tactics) 40–49; *Eulamprus tympanum* (variation in litter size and offspring size) 390–395; *Hyla chrysoscelis* and *H. versicolor* (calling sites in sympatric choruses in Missouri) 373–382; *Hyperolius marmoratus* (male behavior and mating success) 236–246.
SATELLITE BEHAVIOR, *Eleutherodactylus johnstonei* 40–49.
SEX DETERMINATION, 6 spp. of turtle in S Mexico (effects of incubation temperature) 265–270.
SEXUAL DIMORPHISM, *Naja philippinensis* (size of fang discharge orifice) 424–434.
SEXUAL STRATEGY, *Eleutherodactylus johnstonei* 40–49.
SKIN, *Hynobius nigrescens* (effect of prolactin) 279–287; *Sphaerodactylus roosevelti* (extreme mechanical weakness) 124–133.
SONOGRAM, *Eleutherodactylus coqui* (aggressive call) 53.
SOUNDS, *Caretta caretta* 1–30.
SPACING, *Eleutherodactylus johnstonei* 40–49.
SPERMATOZOON, *Cnemidophorus sexlineatus* (ultrastructure) 330–343.
SWIMMING, *Caretta caretta* 1–30.
SYSTEMATICS, *Eleutherodactylus corona* and *E. dolomedes* (new spp. from Haiti); *Lepidophyma* (3 Mexican spp.) and the status of *Gaigeia* 97–110; *Naja* (population systematics, Asiatic) 69–85; *Nerodia* (mtDNA relationships among 6 spp.) 60–68; *Sphaerodactylus* (2 new spp. from E Cuba) 358–367.
TADPOLE, *Eupsophus vertebralis* 115–119; *Hyla allenorum* and *H. fasciata* (with a key to the *H. parviceps* group) 439–447; *Ptychohyla sanctaecrucis* (new sp. from Guatemala) 159.
TAIL BREAKAGE, *Coniophanes fissidens* 448–455.
TAXONOMY, *Arrhyton ridimitum* (status uncertain because of lost holotype) 168–177; *Arrhyton supernum* and *A. procernum* (new spp. from Cuba) 168–177; *Duellmanohyla* (new genus of Central American hylid frogs) 153–167; *Duellmanohyla chamulae* (new combination, formerly *Ptychohyla schmidtorum chamulae*) 153–167; *Eleutherodactylus cacao* (new sp. from SW Colombia) 347–350; *Eleutherodactylus corona* and *E. dolomedes* (new spp. from Haiti) 351–358; *Gonocephalus* (Asian and Australian spp. have strikingly different karyotypes) 120–124; *Hydrophis bituberculatus* (rediscovery and redescription) 85–97; *Lepidophyma* (*Gaigeia*) (3 spp. in Mexico: *L. dontomasi, L. gaigeae,* and *L. radula*) 97–110; *Naja naja* (populations in India and central Asia represent 3 spp., *N. naja, N. kaouthia,* and *N. oxiana*) 69–85; *Nerodia paucimaculata* (distinct from *N. harteri*) 60–68; *Phrynops kempffi* (new sp. of frog from Bolivia) 111–114; *Ptychohyla* (genus redefined to include 10 spp.) 153–167; *Ptychohyla macrotympanum* (elevated to species status, formerly *P. euthysanota*

macrotympanum) 153–167; *Ptychohyla sanctaecrucis* (new sp. from Guatemala) 153–167; *Sphaerodactylus* (2 new spp. from E Cuba) 358–367.
TEMPERATURE, *Xantusia riversiana* 271–279.
TERRITORIALITY, *Eleutherodactylus johnstonei* (Barbados) 40–49.
TESTICULAR CYCLE, *Chelydra serpentina* (in Wisconsin) 193–201.
THERMOREGULATION, *Rana pipiens* and *Sauromalus obesus* (heating and cooling rates compared) 57–60.
TRACKS, *Caretta caretta* 1–30.
VOCALIZATION, *Eleutherodactylus coqui* (aggressive calls) 49–56; *Eleutherodactylus corona* and *E. dolomedes* (new spp. from Haiti) 355.
WATER DRIVE, *Hynobius nigrescens* (induced by prolactin) 279–287.
ZOOGEOGRAPHY, *Gonocephalus* (Asian and Australian spp. have strikingly different karyotypes) 120–124; *Tylototriton andersoni* (allozyme variation among 3 islands of the Ryuku Archipelago related to geological history) 178–184.